

(19) World Intellectual Property Organization
International Bureau




(43) International Publication Date
24 July 2003 (24.07.2003)

PCT


(10) International Publication Number
WO 03/059931 A1

- (51) International Patent Classification⁷: **C07J 51/00**, 1/00, 5/00 (74) Agent: **ROVERS, A.M.A.**; P.O. Box 20, NL-5340 BH Oss (NL).
- (21) International Application Number: PCT/EP03/00339 (81) Designated States (*national*): AE, AG, AL, AU, BA, BB, BR, BZ, CA, CN, CO, CR, CU, DM, DZ, EC, GD, GE, HR, HU, ID, IL, IN, IS, JP, KE, KP, KR, LC, LK, LR, LT, LV, MA, MG, MK, MN, MX, MZ, NO, NZ, PH, PL, RO, RU, SG, TT, UA, US, UZ, VN, YU, ZA.
- (22) International Filing Date: 14 January 2003 (14.01.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 02075230.9 21 January 2002 (21.01.2002) EP (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): **AKZO NOBEL N.V.** [NL/NL]; Velperweg 76, NL-6824 BM Arnhem (NL).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **STOELWINDER, J.** [NL/NL]; Diosynth B.V., P.O. Box 20, NL-5340 BH Oss (NL). **OSTENDORE, M.** [NL/NL]; Diosynth B.V., P.O. Box 20, NL-5340 BH Oss (NL). **BUGGENUM, Van P.A.M.** [NL/NL]; Diosynth B.V., P.O. Box 20, NL-5340 BH Oss (NL).
- Published:**
- with international search report
 - before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: PROCESS FOR THE PREPARATION OF 7 α -METHYLSTEROIDS



(I)



(II)

(57) Abstract: The invention relates to a process for the preparation of 7 α -methyl hydroxy steroids of the Formula I: (Formula I); wherein R₁ is hydrogen, methyl or C \equiv CH; R₂ is (CH₂)_nOH, wherein n is 0, 1 or 2; by a copper mediated 1,6-conjugate addition of a Grignard reagent CH₃MgX, X being a halogen, to the 4,6-unsaturated 3-ketosteroid of formula II: (Formula II); wherein R₁ and R₂ are as previously defined, comprising protecting the hydroxy group of the steroid of Formula II with a trialkylsilyl group, followed by treating the hydroxy protected steroid with the Grignard reagent. The process of the invention is useful for the production of pharmacologically interesting steroids.

PROCESS FOR THE PREPARATION OF 7 α -METHYLSTEROIDS

The invention relates to a new process for the preparation of 7 α -methylsteroids.

5 7 α -Methylsteroids represent an important class of pharmacologically active compounds. One example of a steroid with a methylgroup at the 7-position is (7 α ,17 α)-17-hydroxy-7-methyl-19-norpregn-5(10)-en-20-yn-3-one (Tibolone) which is the active component of Livial®, used as hormone replacement therapy in the treatment of menopausal complaints. Tibolone is a compound exerting tissue-specific hormonal activity. In laboratory and clinical studies, Tibolone
10 exerts estrogenic effects on the thermoregulatory system, vaginal tissue and bone, but does not display estrogenic activity on breast or the endometrium. Tibolone's promising effects on bone are also currently under investigation for the prevention of osteoporosis in women who are likely to develop this condition. (1) Pavlov, P.W., *et al.* Gynecol. Endocrinol. 1999;13: 230–237, 2) Tibolone (Livial). A new steroid for the menopause. Drug Ther. Bull. 1991;29:77-8, 3) Moore
15 R.A. Systematic and economical review for Livial. In: Rymer J (Ed.), Livial: A review of clinical studies. Br.J.Gynaecol. 1999;106 (3 Suppl. 19) 1)

Another pharmacological important class of 7 α -methylsteroids are 7 α -methyl-19-nortestosterone (MENT™) and the related esters at the 17-position (WO99/67271), which are synthetic
20 androgens being developed for hormone replacement therapy and male fertility control. Studies have shown MENT™ to be more potent than the male hormone testosterone in maintaining muscle mass, potency, and libido and in its ability to suppress sperm production. Also, MENT™ does not enlarge the prostate as much as testosterone does, which may result in safer medical use. (1) Ogawa, S., *et al.* 1996; 30:74-84. Hormones and Behavior, 2) Robbins, A., *et al.*
25 Society for Neuroscience Abstracts 1994;20 (part 1):376, 3) Sundaram, K., *et al.* Annals of Medicine 1993;25:199-205, 4) Morali, G. *et al.* Biology of Reproduction 1993;49:577-581).

A third example of a potential interesting steroid with a methylgroup at the 7-position is 3-hydroxy-7 α -methyl-21-[2'-methoxy-4'-(diethylaminomethyl)-phenoxy]-19-norpregna-
1,3,5(10)triene citrate (SR 16234), which is a Selective Estrogen Receptor Modulator (SERM)
30 which has been found to have potent antitumor activity with tissue selective properties, and complete antagonist-antiestrogenic activity in human breast tumor cells. (1) WO 01/58919 A2, 2) WO 99/33859 A2, 3) US 6,281,205 B1, 4) US 20020032180 A1).

From a synthetic point of view, the stereoselective introduction of the substituents at C-7 constitutes the key transformation in the assembly of representatives from this important class of steroids. According to literature, the introduction of an alkyl group in the 7-position of a steroid backbone is generally accomplished by a) cuprous chloride or cupric acetate catalyzed 1,6-conjugate addition of alkylmagnesiumhalogens to 4,6-unsaturated 3-ketosteroids, or b) by conjugate addition of copper-lithiumalkyl-reagents to 4,6-unsaturated 3-ketosteroids. However, in general these methods yield mixtures of 7 α - and 7 β -alkylsteroids (1) Modi, S.P. *et al.*, J. Org. Chem. 1989;54: 2317-2321, 2) Grunwell, J. F. Steroids, 1976;27: 6, 759. 3) Campbell, J.A. *et al.*, J.Am.Chem.Soc. 1959;81: 4069, 4) Grunwell, J.F., *et al.*, Steroids. 1976;27: 750, 5) US patent 3,798,213, 6) van Vliet N.P., *et al.* Recl.Trav.Chim.Pays-Bas 1986;105:111) in α,β -ratios ranging from 1.5:1 to 9:1. Isolation of the pharmacological interesting 7 α -isomers from the accompanying undesirable 7 β -isomers, which are less effective enzyme inhibitors (O'Reilly, J.M. *et al.* J. Med. Chem. 1995;38: 2842) only can be achieved by chromatographic separation, or by laborious work-up procedures by repetitive recrystallization. Both operations decrease the yield of the desired 7 α - isomer significantly.


In WO 01/58919, a 4,6-unsaturated 3-ketosteroid is alkylated at the 7-position with methyllithium in the presence of lithium bromide. In this case it is reported that the stereoselectivity of this 7-alkylation increases after protecting the extant hydroxylgroup with a THP-ether, allegedly by complex formation with the lithium in favor of attack from the α -face of the steroid. Despite the increase in selectivity, unfortunately in this last method the high reactivity of the acetate protecting group with the methyllithium is reflected in only a moderate yield of 67% of the 7 α -methylsteroid.

Alternative methods for the introduction of 7-alkylgroups have been developed, but are not generally applicable. Only a limited selection of alkylgroups can be introduced, the method is limited to the synthesis of steroids with an aromatic A-ring, or additional reaction steps are required, making these methods generally inefficient. For example: 7 α -substituted estradiols have been prepared by conversion of 6-ketoestradiols into 6-(phenylsulfonyl)-6-dehydroestradiol, which undergoes conjugate addition of organolithium reagents to the C-7-position, followed by additional steps to remove the sulfone group. (Künzer, H. *et al.* Tetrahedron Lett. 1994;35: 11, 1691) In a more recent approach 7 α -substituted estradiols have been prepared by alkylation of 6-ketoestradiols, followed by deoxygenation and deprotection

with boron trifluoride etherate and triethylsilane. (Tedesco, R. *et al.* Tetrahedron Lett. 1997;38: 46, 7997). In WO 01/58919, a 6-ketoestradiol is alkylated at the 7-position by reaction with methyl iodide in the presence of lithium diisopropylamine, followed by catalytic removal of the 6-ketone using hydrogen and platinum and palladium; reported is that the stereoselectivity of this 7-alkylation increases after protection of the extant hydroxylgroup with a THP-ether, allegedly by steric hindrance from the β -face of the steroid. In DE 4,018,828 A it is described that 7 α -methylestradiols were prepared by a sequence of reactions from 8,9-unsaturated estradiols with formaldehyde in the presence of Lewis acids, followed by catalytic hydrogenation, tosylation, and reduction. In EP 0,262,201 B1 the preparation of 7 α -propylsteroids is described from 4,6-unsaturated 3-ketosteroids by a Sakurai reaction of allyltrialkylsilanes or allyltrialkyltin compounds in the presence of Lewis acids, followed by a selective hydrogenation with tris(triphenylphosphine)rhodium(I)-chloride (Wilkinson's catalysts). (see also Nickisch, K; Laurent, H. Tetrahedron Lett. 1988;29: 13, 1533).

Therefore, a major challenge in the synthesis of 7-alkylsteroids is control of the diastereoselectivity of the copper mediated 1,6-conjugate addition, which is the most straightforward method of introduction for 7-alkylgroups.


According to the present invention, a process has now been found for the preparation of 7 α -methyl steroids of the formula I



wherein R1 is hydrogen, methyl or $C\equiv CH$,

R2 is $(CH_2)_nOH$, wherein n is 0, 1 or 2;

by a copper mediated 1,6-conjugate addition of a Grignard reagent CH_3MgX , X being a halogen (Cl, Br or I), to the 4,6-unsaturated 3-ketosteroid of formula II,



wherein R1 and R2 are as previously defined, comprising protecting the hydroxy group of the steroid of formula II with a trialkylsilyl group, followed by treating the hydroxy protected steroid with the Grignard reagent.

5 The process of this invention results in a major increase in solubility of the steroid substrate, opens up the possibility to increase the concentration of steroid substrates in said process (a range of concentrations is possible, but preferably the concentration of the steroid is 0.1 to 0.3 molar), and surprisingly shows a markedly improved stereoselectivity in favor of the desired 7 α -isomer. The levels of the unwanted 7 β -isomer are decreased to levels below 2.5%, which is a
10 more than sixfold improvement in selectivity (from approximately 6:1 to 39: 1).

Although the trialkylsilyl protection of a hydroxy group in a process wherein a 7 α -ethyl group was introduced has been described previously (WO 01/05806), the reported selectivity was only 85:15 (α : β).

15 Preferably, R1 is hydrogen, methyl or C \equiv CH and R2 is OH; or R1 is hydrogen and R2 is (CH₂)₂OH. More preferably, when R2 is OH, R1 is hydrogen or C \equiv CH.

The preferred Grignard reagent to be used in the process of the invention is CH₃MgCl.

The Grignard reaction of the process of the invention can be performed in several solvents or
20 mixtures of solvents, which are well known to the person skilled in the art, such as tetrahydrofuran, dimethoxyethane, diethyl ether, mono- and diglyme, toluene and the like. Preferred are tetrahydrofuran or diethyl ether or a mixture of these solvents. Tetrahydrofuran is the most preferred solvent.

The Grignard reagent is used equimolar or in excess to the steroid. Preferably, the molar ratio of
25 the steroid to the Grignard reagent is 1:1 to 1:7.

The step wherein the trialkylsilyl protective group is introduced according to the process of the invention can be integrated in the 1,6-conjugate addition reaction, retaining the improved stereoselectivity in favor of the desired 7 α -isomer and without loss of yield. The term "alkyl" used herein means a (1-4C)alkyl group, being a branched or unbranched alkyl group having 1 to
30 4 carbon atoms, in particular methyl or ethyl. The preferred protective groups are trimethylsilyl or triethylsilyl; most preferred is trimethylsilyl.

In the process of this invention a copper catalyst is used to catalyze the Grignard reaction, such as copper(II) acetate, copper(II) chloride, copper(II) bromide, copper(II) iodide and the like. The preferred catalysts are selected from copper(II) acetate or copper(II) chloride. The most preferred catalyst is copper(II)acetate.

5 The reaction temperature of the Grignard reaction of the process of the invention is not very critical, but should be kept low, preferably between -78 °C and 0 °C. The preferred temperature range is from -35 °C to -25 °C.

This new process for the first time makes a straightforward approach available for increasing the stereoselectivity of copper catalyzed 1,6-conjugate addition of methylmagnesiumhalogens to hydroxylated 4,6-unsaturated 3-ketosteroids. Consequently, laborious work-up procedures by
10 troublesome chromatographic separations or by repetitive recrystallization are unnecessary.

In a suitable process of the invention a hydroxy 4,6-unsaturated 3-ketosteroid is treated with a trialkylsilyl reagent to protect the hydroxy functionality, after which a solution of the silyl
15 protected steroid is added to a mixture of the copper catalyst and the Grignard reagent in an appropriate solvent, which after stirring for some time is followed by removal of the protective group. Removal of the trialkylsilyl protective group takes place under the typical conditions needed for the conjugation to the 4-unsaturated 3-ketosteroids, resulting in an effective one-step procedure. Suitably, the process is finalized by acid treatment for both removal of the trialkylsilyl
20 group and equilibration of the 3,5-unsaturated magnesium-enolate to the desired 7 α -methylsteroids of the formula I.

The process of the invention is particularly useful for the preparation of 7 α -methylsteroid (7 α ,17 α)-17-hydroxy-7-methyl-19-norpregn-4-en-20-yn-3-one, suitable for preparation of
25 (7 α ,17 α)-17-hydroxy-7-methyl-19-norpregn-5(10)-en-20-yn-3-one (Tibolone), for the preparation of 7 α -methyl-19-nortestosterone (MENT), and for the preparation of (7 α)-21-hydroxy-7-methyl-19-norpregn-4-en-3-one, suitable for the preparation of 3-hydroxy-7 α -methyl-21-[2'-methoxy-4'-(diethylaminomethyl)-phenoxy]-19-norpregna-1,3,5(10)triene citrate (SR 16234). For comparison, without introduction of a trialkylsilyl protective group, MENT was
30 prepared by a copper catalyzed 1,6-conjugate addition from 17 β -17-hydroxy-estra-4,6-diene-3-one in approximately 55% yield, with a 7 α , β -ratio of 85:15 (according to procedures described

in US 5,342,834; FR 4.521 M). Isolation of the 7 α -isomer could only be achieved by repeated crystallizations from respectively heptane, and aqueous acetone, decreasing the overall yield to a moderate 44%. In a suitable process according to the invention, introduction of a trimethylsilyl protective group before the 1,6-conjugate addition reaction increases the overall yield of MENT to 79%, which is an improvement of the overall yield by 80%.

A further object of the present invention is the compound 21-hydroxy-19-norpregn-4,6-dien-3-one, which is a suitable intermediate for use in the process of the present invention for the preparation of (7 α)-21-hydroxy-7-methyl-19-norpregn-4-en-3-one, which in turn is suitable for the preparation of 3-hydroxy-7 α -methyl-21-[2'-methoxy-4'-(diethylaminomethyl)-phenoxy]-19-norpregn-1,3,5(10)triene citrate (SR 16234).

The invention is further illustrated by the following examples, which does not mean any limitation.

EXAMPLES

EXAMPLE 1.

7 α -methyl-19-nortestosterone (MENT).

Chlorotrimethylsilane (34.4 ml, 370 mmol) was added dropwise under nitrogen atmosphere at 20°C to a solution of (17 β)-17-hydroxy-estra-4,6-diene-3-one (25 g, 92 mmol) (Wettstein, A. Helv.Chim.Acta, 1940;23: 388) and triethylamine (50 ml, 350 mmol) in anhydrous tetrahydrofuran (250 ml). The resulting mixture was stirred at reflux for 2 h. The reaction mixture was cooled to room temperature, water (125 ml) was added and the mixture was extracted with toluene (250 ml). The combined organic layer was washed with an 10% aqueous sodium chloride solution (125 ml) and evaporated under reduced pressure to yield the 17-trimethylsilyloxy-intermediate as a residual solid, which without isolation was dissolved in anhydrous tetrahydrofuran (250 ml). This solution was added at -30°C in 2 h to a mixture of copper(II) acetate (2.58 g, 12.5 mmol) and a 3M solution of methylmagnesium chloride (61.6 g,

185 mmol) in tetrahydrofuran (120 ml), and the orange-red solution was stirred at -30°C for 1 h. Next the reaction mixture was poured into a solution of sulfuric acid (36.5 ml, 681 mmol) and water (500 ml) and stirred at 40°C for 2.5 h. Sodium acetate was added to a pH of 3.5, and the mixture was extracted with ethyl acetate (260 ml) and washed with aqueous ammonium chloride (83 ml, 10%). After removal of solvents by evaporation under reduced pressure aqueous acetone (150 ml, 50%) was added, and at -10°C 7α -methyl-19-nortestosterone (MENT) was crystallized (19.6 g, 79%) (in accordance to US 5,342,834; FR 4.521 M) in a $7\alpha,\beta$ -ratio of 99:1. $^1\text{H-NMR}$ (400 MHz, CDCl_3): δ 5.68 (bt, 1H, C(4)H), 3.54 (t, 1H, C(17)H), 2.37-0.88 (m, 19H), 0.66 (s, 3H, C(18) H_3), 0.61 (d, 3H, C(7) CH_3).

EXAMPLE 2.

($7\alpha,17\alpha$)-17-Hydroxy-7-methyl-19-norpregn-4-en-20-yn-3-one.

Chlorotrimethylsilane (6.28 ml, 67.5 mmol) was added dropwise under nitrogen atmosphere at 20°C to a solution of (17α)-17-hydroxy-19-norpregna-4,6-dien-20-yn-3-one (5.0 g, 17 mmol) (GB 935116) and pyridine (11.4 ml, 142 mmol) in anhydrous tetrahydrofuran (50 ml). The resulting mixture was stirred at reflux for 2.5 h. The reaction mixture was cooled to room temperature, water (25 ml) was added and the mixture was extracted with toluene (50 ml). The combined organic layer was washed with aqueous methanol (25 ml, 70%) and evaporated under reduced pressure to yield the 17-trimethylsilyloxy-intermediate as a residual solid, which without isolation was dissolved in anhydrous tetrahydrofuran (110 ml). This solution was added at -30°C in 2 h to a mixture of copper(II) acetate (0.58 g, 2.9 mmol) and a 3M solution of methylmagnesium chloride (40 g, 118 mmol) in tetrahydrofuran (50 ml), and was stirred at -30°C for 1 h. Next the reaction mixture was poured into a solution of sulfuric acid (6.8 ml, 126 mmol) and aqueous tetrahydrofuran (160 ml, 15%) and stirred at 20°C for 8 h. Sodium acetate was added to a pH of 3.5, and after removal of solvents by evaporation under reduced pressure ($7\alpha,17\alpha$)-17-hydroxy-7-methyl-19-norpregn-4-en-20-yn-3-one was crystallized (5.2 g, 99%) (in accordance to literature: Van Vliet, N.P., *et al.* Recl.Trav.Chim.Pays-Bas, 1986;105: 111), in a $7\alpha,\beta$ -ratio of 95:5. $^1\text{H-NMR}$ (400 MHz, CDCl_3): δ 5.86 (bt, 1H, C(4)H), 2.59 (s, 1H, C(21)H), 2.52-0.86 (m, 19H), 0.94 (s, 3H, C(18) H_3), 0.79 (d, 3H, C(7) CH_3).

EXAMPLE 3.

21-hydroxy-19-norpregn-4,6-dien-3-one.

4-Toluenesulfonic acid (100 mg, 0.56 mmol) was added under a nitrogen atmosphere at -5 °C to a suspension of 21-hydroxy-19-norpregn-4-en-3-one (30.0 g, 99.2 mmol) (WO 01/58919) and triethyl orthoformate (22.1 mL, 133 mmol) in ethanol (60 mL). After being stirred for 2 h at 0 °C the reaction mixture was quenched with triethylamine (0.22 mL, 1.6 mmol) and heated at reflux for 15 min. The reaction mixture was diluted with ethanol (50 mL), ethyl acetate (1.0 L), water (500 mL) and dichloromethane (500 mL). The organic layer was separated, the aqueous layer was extracted (dichloromethane, 2 x 250 mL) and the combined organic layers were concentrated *in vacuo*. A solution of the residue (50 g) in THF (140 mL) was added to a suspension of chloranil (26 g, 106 mmol) in methanol (140 mL), water (19 mL), acetic acid (5.1 mL) and pyridine (1.9 mL) at 20 °C. After being stirred at this temperature for 2 h, the reaction was quenched with a solution of sodium hydroxide (42 g, 1.05 mol) and sodium hydrosulfite (4.9 g, 28 mmol) in water (490 mL). The aqueous layer was extracted (dichloromethane, 3 x 500 mL), the combined organic layers were washed until pH neutral (water, 4 x 100 mL) and concentrated *in vacuo* to yield 21-hydroxy-19-norpregn-4,6-dien-3-one (25.0 g, 84%). The product was purified by column chromatography and crystallization from ethyl acetate (10.9 g, 37%). ¹H-NMR (400 MHz, CDCl₃): δ 6.25-6.17 (ABX dq, 2H, C(6)H and C(7)H), 5.76 (bd, 1H, C(4)H), 3.73-3.58 (m, 2H, C(21)H₂OH), 2.50-1.08 (m, 19H), 0.69 (s, 3H, C(18)H₃)


(7α)-21-hydroxy-7-methyl-19-norpregn-4-en-3-one.

Chlorotrimethylsilane (2.0 mL, 33.3 mmol) was added dropwise under a nitrogen atmosphere at 20°C to a suspension of 21-hydroxy-19-norpregn-4,6-dien-3-one (5.0 g, 17 mmol) and triethylamine (12 mL, 80 mmol) in anhydrous tetrahydrofuran (100 mL). The resulting mixture was stirred at reflux for 1 h. The reaction mixture was cooled to room temperature, water (50 mL) was added, the organic layer was separated and the water layer was extracted with toluene

(25 mL). The combined organic layers were concentrated *in vacuo*, yielding crude intermediate silylether (11 g). A solution of the silylether in tetrahydrofuran (121 mL) was added in 45 min to a solution of copper acetate (285 mg, 2.86 mmol) and methyl magnesium chloride (3.0 M solution in THF, 38.8 mL, 116 mmol) in THF (45 mL) at -20°C. After being stirred for 1 h at this temperature the reaction was quenched with a solution of sulfuric acid (6.7 g, 68 mmol) in water (125 mL) and THF (25 mL). After being stirred for 17 h at 20°C, the reaction mixture was distilled THF-free and the product was extracted with dichloromethane (3 x 50 mL). After concentration *in vacuo*, (7 α)-21-hydroxy-7-methyl-19-norpregn-4-en-3-one was obtained as an amorphous solid (5.2 g, 99%) (in accordance to literature WO 01/58919) in a 7 α / β ratio of 95:5. ¹H-NMR (400 MHz, CDCl₃): δ 5.85 (bt, 1H, C(4)H), 3.75-3.57 (m, 2H, C(21)H₂OH), 2.50-0.97 (m, 22H), 0.77 (d, 3H, C(7)CH₃), 0.67 (s, 3H, C(18)H₃).

CLAIMS


1. A process for the preparation of 7 α -methyl steroids of the formula I



5 wherein R1 is hydrogen, methyl or C \equiv CH;

R2 is (CH₂)_nOH, wherein n is 0, 1 or 2;

by a copper mediated 1,6-conjugate addition of a Grignard reagent CH₃MgX, X being a halogen, to the 4,6-unsaturated 3-ketosteroid of formula II,



10 wherein R1 and R2 are as previously defined,

comprising protecting the hydroxy group of the steroid of formula II with a trialkylsilyl group, followed by treating the hydroxy protected steroid with the Grignard reagent.

2. The process of claim 1, wherein R1 is hydrogen, methyl or C \equiv CH and R2 is OH; or R1 is
15 hydrogen and R2 is (CH₂)₂OH.

3. The process of claims 1 or 2, wherein the Grignard reagent is CH₃MgCl.

4. The process of any one of claims 1 - 3, wherein the trialkylsilyl group is a trimethylsilyl group.

5. The process of any one of claims 1 - 4, wherein the solvent of the Grignard reaction is
20 tetrahydrofuran, diethyl ether or a mixture thereof.

6. The process of any one of claims 1 - 5, wherein the concentration of the steroid is 0.1 to 0.3 molar.

5 7. The process of any one of claims 1 - 6, wherein the molar ratio of the steroid to the Grignard reagent is 1:1 to 1:7.

8. The process of any one of claims 1 - 7, wherein as copper catalyst copper(II) acetate or copper(II) chloride is used.

10 9. The process of any one of claims 1 - 8, wherein the reaction temperature of the Grignard reaction is -78 °C to 0 °C.

10. The compound 21-hydroxy-19-norpregn-4,6-dien-3-one.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/00339

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07J51/00 C07J1/00 C07J5/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07J

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, BEILSTEIN Data, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 3 341 557 A (BABCOCK JOHN C ET AL) 12 September 1967 (1967-09-12) examples 27,28	1-10
Y	--- CAMPBELL, J. ALLAN ET AL: "7.alpha.-Methyl-19-norsteroids; a new class of potent anabolic and androgenic hormones" STERIODS (1963), 1, 317-24, XP002204811 page 317, last paragraph -page 318, paragraph 1 page 319, scheme, compound IV-Vb --- -/--	1-10

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

23 June 2003

Date of mailing of the international search report

01/07/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Watchorn, P

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/00339

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
L	<p>J. A. CAMPBELL ET AL: "The Synthesis of Some 7.alpha.- and 7.beta.-Methyl Steroid Hormones"</p> <p>JOURNAL OF THE AMERICAN CHEMICAL SOCIETY., vol. 81, 1959, pages 4069-4074, XP002204812</p> <p>AMERICAN CHEMICAL SOCIETY, WASHINGTON, DC., US</p> <p>ISSN: 0002-7863</p> <p>page 4071, column 1, last paragraph -column 2, paragraph 1</p> <p>This document completes the disclosure of the above document with respect to the exact nature of the Grignard reagent used in the 7.-alpha.-methylation of the 19-nor 4,6-dien-3-one compound IV</p> <p>---</p>	1-10
Y	<p>WO 01 05806 A (AKZO NOBEL NV)</p> <p>25 January 2001 (2001-01-25)</p> <p>cited in the application</p> <p>example 1</p> <p>---</p>	1-10
Y	<p>WO 01 40255 A (LEYSEN DIRK ;AKZO NOBEL NV (NL); LOUW JAAP V D (NL); BUMA BURSI RO)</p> <p>7 June 2001 (2001-06-07)</p> <p>example 5</p> <p>---</p>	1-10
Y	<p>WO 00 59920 A (LEYSEN DIRK ;LOUW VAN DER JAAP (NL); AKZO NOBEL NV (NL); BUMA BURS)</p> <p>12 October 2000 (2000-10-12)</p> <p>examples 1,7</p> <p>---</p>	1-10
Y	<p>NICKISCH, KLAUS ET AL: "Stereoselective synthesis of 7.alpha.-allyl- and 7.alpha.-propylsteroids"</p> <p>TETRAHEDRON LETT. (1988), 29(13), 1533-6, XP002204813</p> <p>page 1535, last paragraph</p> <p>---</p>	1-10
Y	<p>WO 01 58919 A (STANFORD RES INST INT)</p> <p>16 August 2001 (2001-08-16)</p> <p>figures 4,8</p> <p>---</p>	1-10
P,Y	<p>WO 02 10188 A (MESTRES JORDI ;LOOZEN HUBERT JAN JOZEF (NL); AKZO NOBEL NV (NL); V)</p> <p>7 February 2002 (2002-02-07)</p> <p>page 8, line 2</p> <p>-----</p>	1-10

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 03/00339

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 3341557	A	12-09-1967	CH 401040 A	31-10-1965
			CH 432508 A	31-03-1967
			CH 421955 A	15-10-1966
			FR 1502605 A	24-11-1967
			GB 941634 A	13-11-1963
WO 0105806	A	25-01-2001	AU 6562900 A	05-02-2001
			BR 0012489 A	02-04-2002
			CA 2379223 A1	25-01-2001
			CN 1360589 T	24-07-2002
			CZ 20020192 A3	12-06-2002
			WO 0105806 A1	25-01-2001
			EP 1203011 A1	08-05-2002
			HU 0201952 A2	28-10-2002
			JP 2003505394 T	12-02-2003
			NO 20020222 A	25-01-2002
			SK 712002 A3	04-06-2002
			TR 200200076 T2	22-04-2002
			US 2003087886 A1	08-05-2003
			US 6313108 B1	06-11-2001
			US 2002022609 A1	21-02-2002
WO 0140255	A	07-06-2001	AU 2838301 A	12-06-2001
			CA 2390226 A1	07-06-2001
			CN 1402735 T	12-03-2003
			CZ 20021894 A3	14-08-2002
			WO 0140255 A2	07-06-2001
			EP 1237904 A2	11-09-2002
			HU 0203611 A2	28-03-2003
			JP 2003515543 T	07-05-2003
			NO 20022564 A	30-05-2002
			SK 7582002 A3	10-09-2002
			US 2003100543 A1	29-05-2003
WO 0059920	A	12-10-2000	AU 5209600 A	23-10-2000
			WO 0059920 A2	12-10-2000
			EP 1212345 A2	12-06-2002
			JP 2002541153 T	03-12-2002
WO 0158919	A	16-08-2001	WO 0158919 A2	16-08-2001
			US 2001039269 A1	08-11-2001
WO 0210188	A	07-02-2002	AU 8201601 A	13-02-2002
			WO 0210188 A1	07-02-2002
			EP 1307471 A1	07-05-2003
			NO 20030419 A	27-01-2003